

Glucose Management Tenets

- 1. Develop and implement an institutional protocol for peri-operative blood sugar management with goal of 120-180 mg/dL.
- 2. Use an insulin infusion for blood sugar management in the intra-op phase and during the first 24 hours postop.
- 3. Check pre-op HA1C levels for all diabetic (at risk/suspected) patients.
- 4. For non-urgent cases in patients with HA1C > 10%, consider delaying surgery and obtaining better glucose control if possible.
- 5. Establish periodic surveillance of compliance of blood sugar target goals.
- 6. Inpatient consult for diabetic education for newly diagnosed patients and those with uncontrolled blood sugars pre-op.
- 7. Social work consult prior to discharge for assistance with obtaining affordable medications at discharge.
- 8. Schedule specialty (Endocrinology) or primary care follow-up at discharge for blood sugar management.

References:

- Association Between Hemoglobin A1c and Major Adverse Coronary Events in Patients with Diabetes Following Coronary Artery Bypass Surgery. Turgeon RD, et al. Pharmacotherapy. 2020. PMID: 31883378 Clinical Trial. <u>https://accpjournals.onlinelibrary.wiley.com/doi/10.1002/phar.2359</u>
- 2. Elevated preoperative hemoglobin A1c level is predictive of adverse events after coronary artery bypass surgery.

Halkos ME, et al. J Thorac Cardiovasc Surg. 2008. PMID: 18805264

- Elevated preoperative hemoglobin A1c level is associated with reduced long-term survival after coronary artery bypass surgery.
 Halkos ME, Lattouf OM, Puskas JD, Kilgo P, Cooper WA, Morris CD, Guyton RA, Thourani VH.Ann Thorac Surg. 2008 Nov;86(5):1431-7. doi: 10.1016/j.athoracsur.2008.06.078.
- Is there a role for HbA1c in predicting mortality and morbidity outcomes after coronary artery bypass graft surgery? Tennyson C, Lee R, Attia R.Interact Cardiovasc Thorac Surg. 2013 Dec;17(6):1000-8. doi: 10.1093/icvts/ivt351. Epub 2013 Sep 10.
- 5. Relationship between preoperative **hemoglobin A1c** levels and long-term mortality after **coronary artery bypass** grafting in patients with type 2 diabetes mellitus.
- 6. Kuhl J, Sartipy U, Eliasson B, Nyström T, Holzmann MJ.Int J Cardiol. 2016 Jan 1;202:291-6. doi: 10.1016/j.ijcard.2015.09.008. Epub 2015 Sep 12.